

#2.

26 Letters 10 integers.

A password of 6 character is randomly selected.

Find the probability that it consists only of letters or exactly 1 case 'm'.

A = consists only of letter

B = exactly 1 'm'.

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\#A = 26^6 \quad \begin{array}{c} 26 \\ \square \end{array} \begin{array}{c} 26 \\ \square \end{array} \begin{array}{c} 26 \\ \square \end{array} \begin{array}{c} 26 \\ \square \end{array} \begin{array}{c} 26 \\ \square \end{array} \begin{array}{c} 26 \\ \square \end{array}$$

$$\#B = 6 \cdot 25^5 \quad \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \quad \begin{array}{c} \text{m} \end{array}$$

$$\#A \cap B = 6 \cdot 25^5$$

$$\begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \begin{array}{c} 25 \\ \square \end{array} \quad \begin{array}{c} \text{m} \end{array}$$

$$\# \text{ total} = 36^6$$

$$P(A \cup B) = \frac{\#A + \#B - \#A \cap B}{\# \text{ total}}$$

$$= 0.2598$$

m	□	□	□	□	□
□	m	□	□	□	□
□	□	m	□	□	□
□	□	□	m	□	□
□	□	□	□	m	□
□	□	□	□	□	m

#11. Computer chips 7% chance of defect.

#11. Computer chips 7% chance of defect.

Suppose we continue to test until we find 10 that have no defects.

X = # of defective chips that when we find 10th good chip.

Find $P(X=5)$

□ □ . . . □ □

Negative Binomial

$$n = x + 10$$

$$r = 10$$

$$p = 0.93$$

($x+10$) chips in total

stop

$$P(X=x) = \binom{x+10-1}{9} 0.93^{10} 0.07^{(x+10)-10} \quad \text{N.B.M.F.}$$

$$x=5.$$

$$= 0.0016285$$

#13. 85 conforming items
25 non-conforming items.

A sample of 10 is chosen.

Find the mean of non-conforming.

Hypergeometric

$$r, k, (N-k)$$

Hypergeometric

$$N = 85 + 25 = 110$$

$$k = 25$$

$$n = 10$$

$$\frac{\binom{K}{10} \binom{N-K}{x-10}}{\binom{N}{x}}$$

$$E(x) = \mu = n \cdot \frac{K}{N} = 10 \cdot \frac{25}{110} = \frac{25}{11}$$

#12. $p = 0.01$ cracking an egg.

Find, on average, how many cartons (containing one dozen eggs) would you have to inspect in order find 3 cartons with no cracked eggs.

$$p' = (1 - 0.01)^{12} \quad \# \text{ no cracked eggs for a carton.}$$

$$r = 3.$$

Negative binomial.

$$E(x) = \mu = \frac{r}{p} = \frac{3}{(1-0.01)^{12}} = 3.3845.$$

Chapter 5:

Two or More Random Variables

Example : 2 dice

X is fair dice.

X is fair dice.

Y is unfair dice

$$P(Y=y) = \begin{cases} \frac{1}{6} & . & 1 \\ \frac{1}{6} & . & 2 \\ \frac{1}{6} & . & 3 \\ \frac{1}{5} & . & 4 \\ \frac{1}{5} & . & 5 \\ \frac{1}{10} & . & 6 \end{cases}$$

X & Y independent

$$P(X=x, Y=y) = \left\{ \begin{array}{ll} \frac{1}{6} \cdot \frac{1}{6} & x=1, y=1 \\ \frac{1}{6} \cdot \frac{1}{6} & x=1, y=2 \\ & \vdots \\ \frac{1}{6} \cdot \frac{1}{10} & x=6, y=6 \end{array} \right\} 36 \text{ cases.}$$

Joint P.M.F of X, Y

The joint probability mass function:

X and Y Discrete R.V.

Then $f_{XY}(x, y) = P(X=x, Y=y)$

(1) $f_{XY}(x, y) \geq 0$

(2) $\sum_x \sum_y f_{XY}(x, y) = 1$

The joint probability density function:

X and Y Continuous R.V.

Then $f_{XY}(x, y)$.

$$(1) f_{XY}(x, y) \geq 0$$

$$(2) \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_{XY}(x, y) dx dy = 1$$

(3) For any region B for two dimensional

$$P(X, Y \in B) = \iint_B f_{XY}(x, y) dx dy$$

Example: X, Y are Continuous:
joint p.d.f:

$$f(x, y) = cx^2 + \frac{xy}{3} \quad 0 \leq x \leq 1, 0 \leq y \leq 2$$

(a) Find (C) .

$$\iint_{-\infty}^{\infty} f(x, y) dx dy = 1$$

$$\int_0^1 \int_0^2 cx^2 + \frac{xy}{3} dy dx = 1$$

$$\int_0^1 \left(cx^2 y + \frac{x}{3} \cdot \frac{1}{2} y^2 \right) \Big|_0^2 dx = 1$$

$$\int_0^1 2cx^2 + \frac{2}{3}x dx = 1.$$

$$\left[C \cdot \frac{2}{3} x^3 + \frac{1}{3} x^2 \right] \Big|_0^1 = \frac{2}{3}C + \frac{1}{3} = 1$$

$$\therefore C = 1$$

$$\therefore C = 1$$


(b) Find $P(X + Y \geq 1)$.

$$P(Y \geq 1 - X) = P(X + Y \geq 1)$$

$$X \in [0, 1] \quad 1 - X \in [0, 1]$$

$$Y \in [0, 2] \quad Y \geq 1 - X$$

$$\Rightarrow Y \in [1 - X, 2]$$



$$P(X + Y \geq 1)$$

$$= P(Y \geq 1 - X)$$

$$= \int_0^1 \int_{1-x}^2 \left(x^2 + \frac{xy}{3}\right) dy dx$$

$$= \int_0^1 \left(x^2 y + \frac{x}{6} y^2\right) \Big|_{1-x}^2 dx$$

$$= \int_0^1 \left(\frac{4}{3} x^2 + \frac{1}{2} x + \frac{5}{6} x^3\right) dx$$

$$= \left(\frac{4}{9} x^3 + \frac{1}{4} x^2 + \frac{5}{24} x^4\right) \Big|_0^1$$

$$= \frac{65}{72}$$

Example:

X : time until a computer server connects to your machine.

connects to your machine.

Y : time until the server authorizes you as a valid user.

$$f_{X,Y}(x,y) = 6 \cdot 10^{-6} \exp(-0.001x - 0.002y)$$

$$\text{for } \boxed{x < y} \quad \begin{cases} x > 0 \\ y > 0 \end{cases}$$

$$\text{Find } P(X \leq 1000, Y \leq 2000)$$

$$P(X \leq 1000, Y \leq 2000)$$

$$= \int_0^{1000} \int_{\boxed{x}}^{2000} 6 \cdot 10^{-6} \exp(-0.001x - 0.002y) dy dx$$

$$= 6 \cdot 10^{-6} \cdot \int_0^{1000} \left[\int_x^{2000} \exp(-0.002y) dy \right] \exp(-0.001x) dx$$

$$= 6 \cdot 10^{-6} \int_0^{1000} \left[\frac{\exp(-0.002y)}{-0.002} \Big|_x^{2000} \right] \exp(-0.001x) dx$$

$$= 6 \cdot 10^{-6} \int_0^{1000} \frac{\exp(-0.003x) - \exp(-4) \exp(-0.001x)}{0.002} dx$$

$$= 0.003 \left[\frac{\exp(-0.003x)}{0.003} - \frac{\exp(-4) \exp(-0.001x)}{-0.001} \right] \Big|_0^{1000}$$

$$= 1 - e^{-3} - 3(e^{-4} - e^{-5})$$

Marginal probability distribution:

$$f_{XY}(x, y) \Rightarrow \begin{matrix} f_X(x) & (f_X) \\ f_Y(y) & (f_Y) \end{matrix}$$

Discrete one:

$$f_X(x) = \sum_y f_{XY}(x, y)$$

$$f_Y(y) = \sum_x f_{XY}(x, y)$$

Continuous

$$\begin{aligned} f_X(x) &= \int f_{XY}(x, y) dy \\ f_Y(y) &= \int f_{XY}(x, y) dx \end{aligned}$$

} Integral over points in the range.

Example:

$$P(Y > 2000)$$

$$f_{XY}(x, y) = 6 \cdot 10^{-6} e^{-0.001x - 0.002y}, \quad x < y$$

$$f(y) = \int_0^y f_{XY}(x, y) dx$$

$$f(y) = \int_0^y f_{XY}(x, y) dx$$

$$= \int_0^y 6 \cdot 10^{-6} \cdot e^{-0.001x - 0.002y} dx$$

$$= 6 \cdot 10^{-3} e^{-0.002y} [1 - e^{-0.001y}] \text{, for } y > 0$$

$$P(Y > 2000) = \int_{2000}^{\infty} f(y) dy$$

$$= \int_{2000}^{\infty} 6 \cdot 10^{-3} [e^{-0.002y} - e^{-0.003y}] dy$$

$$= 6 \cdot 10^{-3} \left[\frac{e^{-0.002y}}{-0.002} - \frac{e^{-0.003y}}{-0.003} \right] \Bigg|_{2000}^{\infty}$$

$$= 0.05$$

$$P(Y > 2000) = \int_{2000}^{\infty} \int_0^y f_{XY}(x, y) dx dy$$

Conditional Distribution and Independent.

X, Y . Discrete R.V.

$$P_{Y|X}(y|x) = \frac{P(X=x \text{ \& } Y=y)}{P(X=x)} \quad P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Properties:

$$(1) P_{Y|X}(y|x) \geq 0$$

$$(2) \sum P_{Y|X}(y|x) = 1$$

$$\sum_y P(Y=y|X=x) = 1$$

$$(3) P(Y \in B | X=x) = \sum_B f_{Y|X}(y|x)$$

for any region B

Continuous X, Y R.V.

$$f_{Y|X}(y|x) = \frac{f_{XY}(x, y)}{f_X(x)} \quad P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Properties:

$$(1) f_{Y|X}(y|x) \geq 0$$

$$(2) \int f_{Y|X}(y|x) dy = 1$$

$$(3) P(Y \in B | X=x) = \int_B f_{Y|X}(y|x) dy$$

for any region B.

Independent:

$$\begin{aligned} P(A \cap B) &= P(A) \cdot P(B) \\ P(A|B) &= P(A) \end{aligned}$$

Discrete

$$(1) P(X=x, Y=y) = P(X=x) \cdot P(Y=y)$$

for all x, y .

$$(2) P_{Y|X}(y|x=x) = P(Y)$$

Continuous:

$$(1) f_{XY}(x, y) = f_X(x) \cdot f_Y(y)$$

$$\dots$$

$$(2) f_{Y|X}(y|x) = f_Y(y)$$

Example:

R.V. x, y with joint p.m.f:

	$Y=2$	$Y=4$	$Y=5$	
$X=1$	$\frac{1}{12}$	$\frac{1}{24}$	$\frac{1}{24}$	$\frac{1}{6}$
$X=2$	$\frac{1}{6}$	$\frac{1}{12}$	$\frac{1}{8}$	$\frac{3}{8}$
$X=3$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{12}$	$\frac{1}{4}$
	$\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{4}$	

$$(c) (X|Y=2)$$

$$P(X=1|Y=2) = \frac{\frac{1}{12}}{\frac{1}{2}} = \frac{1}{6}$$

$$= \frac{P(X=1, Y=2)}{P(Y=2)}$$

(a) Find $P(X \leq 2, Y \leq 4)$

$$P(X \leq 2, Y \leq 4) = \frac{3}{8}$$

(b) Find Marginal distribution for x, y .

$$P(Y=y) = \begin{cases} \frac{1}{2} & Y=2 \\ \frac{1}{4} & Y=4 \\ \frac{1}{4} & Y=5 \end{cases}$$

$$P(X=x) = \begin{cases} \frac{1}{6} & x=1 \\ \frac{3}{8} & x=2 \\ \frac{1}{4} & x=3 \end{cases}$$

Example:

joint pdf.

Example:

joint p.d.f.

$$f(x,y) = cxy \quad 0 \leq x \leq 1, \quad 0 \leq y \leq \sqrt{x}$$

(a) Find 'c'.

$$\iint f(x,y) dx dy = 1$$

$$\int_0^1 \int_0^{\sqrt{x}} cxy dy dx = 1$$

$$\int_0^1 \left(\frac{c}{2} x y^2 \Big|_0^{\sqrt{x}} \right) dx = 1$$

$$\int_0^1 \frac{c}{2} x^2 dx = 1$$

$$\frac{c}{6} x^3 \Big|_0^1 = 1$$

$$\therefore c = 6$$

(b) Find $f_X(x)$ and $f_Y(y)$.

$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x,y) dy$$

$$= \int_0^{\sqrt{x}} 6xy dy$$

$$= 3x^2, \quad 0 \leq x \leq 1$$

$$f_Y(y) = \int_{-\infty}^{\infty} f_{XY}(x,y) dx$$

$$= \int_{y^2}^1 6xy dx$$

$$0 \leq y \leq \sqrt{x}$$

$$\Leftrightarrow y^2 \leq x \leq 1$$

$$= \int_{y^2}^1 6xy \, dx$$

$$\rightarrow y = \text{constant}$$

$$= 3y(1-y^4) \quad . \quad 0 \leq y \leq 1$$

(c) Find $f_{X|Y}(x)$

$$f_{X|Y}(x) = \frac{f_{XY}(x, y)}{f_Y(y)} \quad \boxed{y \text{ is given.}}$$

$$= \boxed{\frac{6xy}{3y(1-y^4)} \quad , \quad y^2 \leq x \leq 1}$$

(d) Are X, Y independent.

$$f_{X|Y}(x) \neq f_X(x)$$

$\therefore X, Y$ are dependent.

(e) $E(X|Y=y)$

$$E(X|Y=y) = \int x \cdot f_{X|Y}(x|y) \, dx$$

$$E(X|Y=y) = \int_{y^2}^1 x \cdot \frac{6xy}{3y(1-y^4)} \, dx$$

$$= \int_{y^2}^1 \frac{2x^2}{1-y^4} \, dx$$

$$= \frac{2}{3} \cdot \frac{1}{1-y^4} \cdot x^3 \Big|_{y^2}^1$$

$$= \frac{2}{3} \frac{1-y^6}{1-y^4} \quad y \text{ is given}$$

$$r = \frac{2}{3} \frac{1-y}{1-y^4} \quad y \text{ is given}$$

$$E(X|Y=2) = \frac{2}{3} \frac{1-2^6}{1-2^4}$$

$$E(X^2|Y=y) = \frac{1-y^8}{2(1-y^4)}$$

$$V(\underline{X}|Y=y) = E(\underline{X^2}|Y=y) - [E(\underline{X}|Y=y)]^2$$

More General:

Suppose we have $X_1, \dots, X_p$ R.V.
 p dimensional.

Joint p.d.f.

$$(1) f_{X_1, \dots, X_p}(x_1, x_2, \dots, x_p) \geq 0$$

$$(2) \underbrace{\int \int \int \dots \int}_{\#p} f_{X_1, \dots, X_p}(x_1, \dots, x_p) \underbrace{dx_1 \dots dx_p}_{\#p} = 1$$

(3) For any Region B .

$$P[(X_1, \dots, X_p) \in B] = \int \int \dots \int_B f_{X_1, \dots, X_p}(x_1, \dots, x_p) dx_1 \dots dx_p$$

Conditional:

$$f(x_1, x_2, \dots, x_5)$$

$$f(x_1, x_2, \dots, x_5)$$

$$f(x_1, x_2 \mid \underline{x_3, x_4, x_5}) = \frac{f(x_1, \dots, x_5)}{\underline{f(x_3, x_4, x_5)}}$$

Independent

$$f_{x_1, \dots, x_p}(x_1, \dots, x_p) = f_{x_1}(x_1) \cdot f_{x_2}(x_2) \cdot \dots \cdot f_{x_p}(x_p)$$

5.2 Covariance and Correlation

X, Y .

The covariance between R.V. X, Y .

denoted by $\text{Cov}(X, Y)$ or σ_{xy}

$$\begin{aligned} \sigma_{xy} = \text{Cov}(X, Y) &= E[(X - \mu_X)(Y - \mu_Y)] \\ &= \underline{E(XY) - E(X)E(Y)} \end{aligned}$$

$$E[h(x, y)] = \begin{cases} \sum \sum h(x, y) f_{XY}(x, y) & (\text{Discrete}) \\ \iint h(x, y) f_{XY}(x, y) dx dy & (\text{Continuous}) \end{cases}$$

Correlation:

ρ_{xy}


$\text{Cov}(X, Y)$

$\sigma_X \sigma_Y$


$$\rho_{xy} = \frac{\text{cov}(X, Y)}{\sqrt{V(X) V(Y)}} = \frac{\sigma_{xy}}{\sigma_x \sigma_y}$$

$$-1 \leq \rho_{xy} \leq 1$$

(1) $\rho_{xy} = 1$ x, y strictly on a line
 $\rho_{xy} = -1$ x, y strictly on a line



which perfectly
linearly correlated



Correlation only measures
 linear relationship

$$\rho_{xy} = 0 = \text{cov}(X, Y)$$

$\Leftrightarrow X, Y$ are independent.

$1 > \rho_{xy} > 0$ positive linearly correlated

$0 > \rho_{xy} > -1$ negative linearly correlated.

Example:
 joint p.d.f.

Example joint p.d.f.

$$f(x, y) = \frac{1}{16} xy, \quad \begin{matrix} 0 \leq x \leq 2 \\ 0 \leq y \leq 4 \end{matrix}$$

Find $\text{Cov}(X, Y)$ and ρ_{xy} .

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y).$$

$$E(XY) = \int_0^4 \int_0^2 xy \cdot f_{XY}(x, y) dx dy$$

$$= \int_0^4 \int_0^2 \frac{1}{16} x^2 y^2 dx dy$$

$$= \frac{1}{16} \int_0^4 \left[\frac{1}{3} x^3 y^2 \Big|_0^2 \right] dy$$


$$= \frac{1}{16} \int_0^4 \frac{8}{3} y^2 dy$$

$$= \frac{1}{16} \frac{8}{9} y^3 \Big|_0^4$$

$$= \frac{32}{9}$$

$$E(X) = \int_0^4 \int_0^2 x \cdot f_{XY}(x, y) dx dy$$

$E(X) = \int x f_X(x) dx$
 $f_X(x) = \int f_{XY}(x, y) dy$



$$f_X(x) = \int_{-\infty}^{\infty} f_{XY}(x, y) dy$$

$$\Rightarrow E(X) = \iint x f_{XY}(x, y) dx dy$$

$$= \frac{1}{16} \int_0^4 \int_0^2 x^2 y dx dy$$

$$= \frac{1}{16} \int_0^4 \frac{8}{3} y dy$$

$$= \frac{4}{3}$$

$$E(Y) = \int_0^4 \int_0^2 y f_{XY}(x, y) dx dy$$

$$= \frac{1}{16} \int_0^4 y^2 \cdot \frac{1}{2} x^2 \Big|_0^2 dy$$

$$= \frac{1}{8} \left(\frac{1}{3} y^3 \Big|_0^4 \right)$$

$$= \frac{8}{3}$$

$$\text{cor}(X, Y) = E(XY) - E(X) \cdot E(Y)$$

$$= 0$$

$\therefore X$ and Y are independent.

5.4 Linear Function of R.V.

shoppers 10p/hour.
24 hours

□ □ □ ... — □
 x_1 x_2 x_3 ... — x_{24}

$$\begin{array}{ccccccc}
 x_1 & x_2 & x_3 & \dots & & & x_{24} \\
 \vdots & \vdots & \vdots & & & & \vdots \\
 & & & & \text{Identical and Independent. I.I.D} & &
 \end{array}$$

$$\begin{aligned}
 E(x_1 + x_2 + \dots + x_{24}) &= E(x_1) + E(x_2) + \dots + E(x_{24}) \\
 &= 24 \cdot E(x) \\
 &= 24 \cdot 10 \\
 &= 240.
 \end{aligned}$$

Linear Combination.

Given R.V. $x_1, x_2, \dots, x_p$ and constants $c_1, c_2, \dots, c_p$.

$$Y = c_1 x_1 + c_2 x_2 + \dots + c_p x_p.$$

is a linear combination of $x_1, \dots, x_p$.

$$E(Y) = c_1 E(x_1) + c_2 E(x_2) + \dots + c_p E(x_p)$$

$$\left[c_1 \int x_1 f(x_1) dx_1 + c_2 \int x_2 f(x_2) dx_2 + \dots \right]$$

$$V(Y) = c_1^2 V(x_1) + c_2^2 V(x_2) + \dots + c_p^2 V(x_p)$$

$$+ \sum_{i < j} \sum c_i c_j \text{Cov}(x_i, x_j)$$

Independent
 $\Leftrightarrow \text{Cov}(x_i, x_j) = 0$

Example: $Y = \underline{2}x_1 + \underline{3}x_2 + 4x_3$

$$\therefore E(Y) = 2E(x_1) + 3E(x_2) + 4E(x_3)$$

$$V(Y) = 4V(x_1) + 9V(x_2) + 16V(x_3)$$

$$V(Y) = 4V(X_1) + 9V(X_2) + 16V(X_3) \\ + 2 \cdot 3 \operatorname{Cor}(X_1, X_2) + 2 \cdot 4 \operatorname{Cor}(X_1, X_3) \\ + 3 \cdot 4 \operatorname{Cor}(X_2, X_3)$$

Example: Geometric and Negative Binomial.

✓ Experiments until one success.

Negative Binomial

↓
Experiments until r success.

$X_1, \dots, X_p \sim \text{Geometric.}$

$Y = X_1 + \dots + X_p \sim \text{Negative Binomial}$
($r = p$)

$X_1, \dots, X_p$ they I. I. D.

identical & independent
distributed.

Hard to prove they are equal

could see $E(X), E(Y)$
 $V(X), V(Y)$

$$E(Y) = \underline{E(X_1)} + \underline{E(X_2)} + \dots + \underline{E(X_r)} \\ = \frac{1}{p} + \frac{1}{p} + \dots + \frac{1}{p}$$

$$= \frac{1}{p}$$

$$V(Y) = 1^2 \cdot V(X_1) + 1^2 \cdot V(X_2) + \dots + V(X_r)$$

$$= \frac{1-p}{p^2} + \dots + \frac{1-p}{p^2}$$

$$= \frac{r(1-p)}{p^2}$$

Example:

Shoppers sell 10 boxes of pop on average per day.

X denote per hour they sell # of boxes.

X follows Normal with $\mu=10$, $\sigma=5$.

(a) what's the probability that 20 days per day in total we sell more than 300 boxes?

$X_1 \dots X_{20}$
 $\uparrow$ 1st Day $\uparrow$ 20th Day.

$$Y = X_1 + \dots + X_{20}$$

$$P(Y \geq 300) ?$$

Standardizing

$$E(Y) = \mu = E(X_1) + \dots + E(X_{20}) = 10 \cdot 20 = 200$$

$$V(Y) = \sigma^2 = 20 \cdot V(X) = 20 \cdot 5^2 = 500.$$

$$V(Y) = \sigma^2 = 20 \quad V(X) = 20 \cdot 5^2 = 500.$$

$$\sigma = \sqrt{500}$$

$$P(Y \geq 300) = P\left(\frac{Y - \mu}{\sigma} \geq \frac{300 - 200}{\sqrt{500}}\right)$$

$$= P(Z \geq 4.47)$$

Symmetric

$$= P(Z \leq -4.47)$$

$$= 0$$

Example:

20 Days.

$X_1, \dots, X_{20} \sim \text{Normal}(\mu=10, \sigma=5).$

What's the probability that on average more than 9 boxes are sold per day.

$$\bar{X} = \frac{X_1 + \dots + X_{20}}{20}$$

$$E(\bar{X}) = \frac{1}{20} E(X_1) + \frac{1}{20} E(X_2) + \dots + \frac{1}{20} E(X_{20})$$

$$= \frac{20}{20} E(X) = E(X) = 10$$

$$V(\bar{X}) = V\left(\frac{1}{20} X_1 + \frac{1}{20} X_2 + \dots + \frac{1}{20} X_{20}\right)$$

$$= \frac{1}{400} V(X_1) + \frac{1}{400} V(X_2) + \dots + \frac{1}{400} V(X_{20})$$

$$= \underbrace{\frac{1}{400} V(X_1) + \frac{1}{400} V(X_2) + \dots + \frac{1}{400} V(X_{20})}_{20}$$

$$= \frac{20}{400} V(X)$$

$$= \frac{1}{20} \cdot 5^2$$

$$= \frac{5}{4}$$

$$P(\bar{X} > 9) = P\left(Z > \frac{9-10}{\sqrt{\frac{5}{4}}}\right)$$

$$= P(Z > -0.89)$$

$$= P(Z < 0.89)$$

$$= 0.8133$$

Example:

X_1, X_2

X_1, X_2 are independent identical.

$$f(x) = x^2(2x + \frac{3}{2}), \quad 0 < x < 1$$

Find $E(Y)$ where $Y = \frac{1}{X_1} + \frac{1}{X_2} + 3$.

$$E(Y) = E\left(\frac{1}{X_1}\right) + E\left(\frac{1}{X_2}\right) + 3$$

$$\neq \frac{1}{E(X_1)}$$

non-linear

$$E(h(x)) = \int h(x) f(x) dx$$

$$E\left(\frac{1}{x}\right) = \int_0^1 \frac{1}{x} \cdot f(x) dx \quad E\left(\frac{1}{x}\right) \neq \frac{1}{E(x)}$$

$$= \int_0^1 \frac{1}{x} \cdot x^2 \left(2x + \frac{3}{2}\right) dx$$

$$= x^2 + \frac{3}{2}x \Big|_0^1$$

$$= \frac{5}{2}$$

$$\therefore E(Y) = \frac{5}{2} + \frac{5}{2} + 3 = 8$$